

# Draft genome sequence of an endophytic bacterium, *Paenibacillus tyrfis* strain SUK123, isolated from *Santiria apiculata* stem



Emmanuel Haruna<sup>a,b</sup>, Noraziah M. Zin<sup>a,\*</sup>, Jonathan M. Adams<sup>c,\*</sup>

<sup>a</sup> Programme of Biomedical Science, School of Diagnostic and Applied Health Sciences, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, Kuala Lumpur, Malaysia

<sup>b</sup> Department of Biochemistry Kaduna State University, Kaduna, Nigeria

<sup>c</sup> Division of Agrifood and Environment, School of Water, Energy and Environment, Building 52a, Cranfield University, Cranfield, Bedfordshire MK43 0AL, UK

## ABSTRACT

Here we report the draft genome sequence of an endophytic *Paenibacillus tyrfis* strain isolated from the Universiti Kebangsaan Malaysia reserve forest, Malaysia. The genome size was approximately 8.04 Mb, and the assembly consisted of 107 scaffolds with 168 contigs, and had a G + C content of 53%. Phylogenetic analysis of strain SUK123 using the 16S rRNA gene revealed that it belonged to the family *Paenibacillaceae* with the highest similarity to *Paenibacillus elgii* SD<sup>T</sup> (99%). Whole genome comparison of SUK123 with related species using average nucleotide identity (ANI) analysis revealed a similarity of 98% to *Paenibacillus tyrfis* Mst1<sup>T</sup>, 94% to *Paenibacillus elgii* B69<sup>T</sup>, 91% to *Paenibacillus ehimensis* A2<sup>T</sup>, 68% to *Paenibacillus polymyxa* SC2<sup>T</sup> and 69% to *Paenibacillus alvei* DMS29<sup>T</sup>. The draft genome was deposited at the European Nucleotide Archive (PRJEB21373).

## Specifications

Organism/cell line/tissue	<i>Paenibacillus tyrfis</i>
Sex	Not applicable
Sequencer or array type	Illumina Miseq
Data format	Raw data and analyzed i.e. assembled
Experimental factors	<i>Paenibacillus tyrfis</i> strain was isolated from <i>Santiria apiculata</i> stem
Experimental features	Isolation of bacteria, Genome sequencing, de novo assembly
Consent	Not applicable
Sample source location	Universiti Kebangsaan Malaysia reserve forest, Latitude & Longitude – 2.9125 & 101.7872

## 1. Direct link to deposited data

<http://www.ebi.ac.uk/ena/data/view/PRJEB21373>.

## 2. Introduction

The bacterial genus *Paenibacillus* has been isolated from many environments, mostly from plant organs and their surrounding soil [1]. The species of *Paenibacillus* was included in the genus *Bacillus* until 1993 when it was proposed that the member of the “16S rRNA group 3” bacilli be distinguished from it [2]. Presently, the genus *Paenibacillus* consists of 395 known species [3]. Most members of the genus *Paenibacillus* are gram variable, either aerobic or facultatively anaerobic, rod-shaped, and endospore-forming with peritrichous flagella for motility [4]. The DNA G + C content of the various species of *Paenibacillus* ranges between 39 and 54 mol% whilst anteiso-C<sub>15:0</sub> is the major cellular fatty acid and meso-diaminopimelic acid is the cell wall peptidoglycan diamino acid [5]. Most members of this genus have been reported to be producers of either active antimicrobial metabolites such as lipopeptides [6,7], plant-growth promoting hormones [8,9], or insecticides [10].

## 3. Experimental design, materials and methods

The endophytic *Paenibacillus tyrfis*, strain SUK123, was isolated from

\* Corresponding authors.

E-mail addresses: [noraziah.zin@ukm.edu.my](mailto:noraziah.zin@ukm.edu.my) (N.M. Zin), [j.m.adams@cranfield.ac.uk](mailto:j.m.adams@cranfield.ac.uk) (J.M. Adams).

**Table 1**  
Statistics of assembled sequence length.

No. of all scaffolds	107
Bases in all scaffolds	8,041,385
No. of large scaffolds (> 1000 bp)	73
Bases in large scaffolds	8,024,609
Largest length	505,699
Scaffold N50	257,089
Scaffold N90	62,404
G + C content (%)	53.061
No. of all contigs	168
Bases in all contigs	8,041,324
No. of large contigs (> 1000 bp)	114
Bases in large contigs	8,013,898
Largest length	501,150
Contig N50	197,909
Contig N90	48,401

the *Santiria apiculata* stem located at the Universiti Kebangsaan Malaysia forest reserve, whilst screening for endophytes with antimicrobial potential against ESKAPE (*Enterococcus faecium*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Acinetobacter baumannii*, *Pseudomonas aeruginosa*, and *Enterobacter species*) pathogens. After sample collection, the stem surface was sterilized to avoid surface-bound epiphytes as described previously [11]. The sterilized stem was cut into a 2 cm section and plated on water agar plate medium from where SUK123 (an antibiotic producing bacterium) was isolated.

Endophytic *Paenibacillus tyrfis* DNA was extracted using an Ultraclean Microbial DNA Isolation Kit as described by the manufacturer (Mo Bio Laboratories, 2746 Loker Ave W # A, Carlsbad, CA 92010, USA). The bacterium was identified by sequencing its 16S rRNA gene. Sequence was performed on Illumina Miseq platform (Majorbio, China) by 2 × 400 bp paired-end libraries. The raw sequence quality was assessed using PRINSEQ lite version 0.20.4 and the genome was assembled using SOAPdenovo v2.04 with all parameters set by default [12] and GapCloser v1.12 was used to fill local inner gaps. The open reading frames (ORFs) were predicted using Glimmer 3.02 [13]. The

biological functions of these predicted ORFs were annotated using various databases.

#### 4. Data description

The analyses of the assembled genome revealed a genome size of about 8,041,385 bp made of 107 scaffolds, 168 contigs, and with a G + C content of 53.06% and N50 contigs size of 197,909 (Table 1). The genome was annotated using Rapid Annotation Subsystems Technology server [14].

The RAST server prediction revealed 7368 coding sequences (CDS) with a total of 2880 CDS (40%) constituting 2747 and 133 of non-hypothetical and hypothetical proteins respectively in the subsystem coverage. A total of 4488 CDS (60%) comprising 2003 and 2485 of non-hypothetical and hypothetical proteins respectively were outside the subsystem coverage (Fig. 1).

The average nucleotide identity (ANI) [15] analysis of SUK123 with closely related species, revealed a similarity index with *Paenibacillus tyrfis* Mst1<sup>T</sup> (98.06%), *Paenibacillus elgii* B69<sup>T</sup> (93.98%), *Paenibacillus ehimensis* A2<sup>T</sup> (91.17%), *Paenibacillus alvei* DMS29<sup>T</sup> (68.81%) and *Paenibacillus polymyxa* SC2<sup>T</sup> (68.27%), suggesting our isolate most closely related to *Paenibacillus tyrfis* Mst1<sup>T</sup>.

#### Declaration of interest

We declare no conflict of interest with respect to this article titled ‘Draft genome sequence of an endophytic *Paenibacillus tyrfis* strain SUK123 isolated from *Santiria apiculata* stem’ submitted to Genomics data journal.

#### Acknowledgement

We appreciate University Kebangsaan University Forest Reserve and Herbarium staff and grateful to Muhanna Al-Shabanni, Aishah Ismail, Radhiah Binti Khairon, and Nur Faizah Abu Bakar, for their assistance during sampling. This work was funded by the Universiti Kebangsaan Malaysia Research Grant (GUP-2015-042).




Fig. 1. A diagrammatic representation of *Paenibacillus tyrfis* strain genome subsystem coverage as annotated by RAST.

## References

- [1] E.N. Grady, J. MacDonald, L. Liu, A. Richman, Z.-C. Yuan, E. Novelli, Current knowledge and perspectives of *Paenibacillus*: a review, *Microb. Cell Factories* 15 (2016) 203, <http://dx.doi.org/10.1186/s12934-016-0603-7>.
- [2] C. Ash, F.G. Priest, M.D. Collins, Molecular identification of rRNA group 3 bacilli (Ash, Farrow, Wallbanks and Collins) using a PCR probe test. Proposal for the creation of a new genus *Paenibacillus*, *Antonie Van Leeuwenhoek* 64 (1993) 253–260.
- [3] S.-H. Yoon, S.-M. Ha, S. Kwon, J. Lim, Y. Kim, H. Seo, J. Chun, Introducing EzBioCloud: a taxonomically united database of 16S rRNA gene sequences and whole-genome assemblies, *Int. J. Syst. Evol. Microbiol.* 67 (2017) 1613–1617, <http://dx.doi.org/10.1099/ijsem.0.001755>.
- [4] Y.-K. Aw, K.-S. Ong, L.-H. Lee, Y.-L. Cheow, C.M. Yule, S.-M. Lee, Newly isolated *Paenibacillus tyrfis* sp. nov., from Malaysian tropical peat swamp soil with broad spectrum antimicrobial activity, *Front. Microbiol.* 7 (2016) 219, <http://dx.doi.org/10.3389/fmicb.2016.00219>.
- [5] O. Shida, H. Takagi, K. Kadowaki, L.K. Nakamura, K. Komagata, Transfer of *Bacillus alginolyticus*, *Bacillus chondroitinus*, *Bacillus curdlanolyticus*, *Bacillus glucanolyticus*, *Bacillus kobensis*, and *Bacillus thiaminolyticus* to the genus *Paenibacillus* and emended description of the genus *Paenibacillus*, *Int. J. Syst. Bacteriol.* 47 (1997) 289–298, <http://dx.doi.org/10.1099/00207713-47-2-289>.
- [6] Z. Huang, Y. Hu, L. Shou, M. Song, Isolation and partial characterization of cyclic lipopeptide antibiotics produced by *Paenibacillus ehimensis* B7, *BMC Microbiol.* 13 (2013) 87, <http://dx.doi.org/10.1186/1471-2180-13-87>.
- [7] G. Aktuganov, A. Melentjev, N. Galimzianova, E. Khalikova, T. Korpela, P. Susi, Wide-range antifungal antagonism of *Paenibacillus ehimensis* IB-X-b and its dependence on chitinase and beta-1,3-glucanase production, *Can. J. Microbiol.* 54 (2008) 577–587, <http://dx.doi.org/10.1139/w08-043>.
- [8] K. Zhou, M. Yamagishi, M. Osaki, *Paenibacillus* BRF-1 has biocontrol ability against *Phialophora gregata* disease and promotes soybean growth, *Soil Sci. Plant Nutr.* 54 (2008) 870–875, <http://dx.doi.org/10.1111/j.1747-0765.2008.00308.x>.
- [9] B. Pichard, D. Thouvenot, Effect of *Bacillus polymyxa* seed treatments on control of black-rot and damping-off of cauliflower, *Seed Sci. Technol.* 27 (1999) 445–465.
- [10] S. Neung, X. Nguyen, K. Naing, Y. Lee, Insecticidal potential of *Paenibacillus elgii* HOA73 and its combination with organic sulfur pesticide on diamondback moth, *Plutella xylostella*, *Appl. Biol.* 55 (2014) 181–186.
- [11] E. Haruna, N.M. Zin, D. Kerfahi, J.M. Adams, Extensive overlap of tropical rain-forest bacterial endophytes between soil, plant parts, and plant species, *Microb. Ecol.* (2017) 1–16, <http://dx.doi.org/10.1007/s00248-017-1002-2>.
- [12] R. Luo, B. Liu, Y. Xie, Z. Li, W. Huang, SOAPdenovo2: an empirically improved memory-efficient short-read de novo assembler, *Gigascience* 1 (2012) 18.
- [13] A.L. Delcher, K.A. Bratke, E.C. Powers, S.L. Salzberg, Identifying Bacterial Genes and Endosymbiont DNA With Glimmer, 23 (2007), pp. 673–679, <http://dx.doi.org/10.1093/bioinformatics/btm009>.
- [14] R. Aziz, D. Bartels, A. Best, The RAST Server: rapid annotations using subsystems technology, *BMC* 9 (2008) 75.
- [15] S.-H. Yoon, S. Ha, J. Lim, S. Kwon, J. Chun, A large-scale evaluation of algorithms to calculate average nucleotide identity, *Antonie Van Leeuwenhoek* (2017) 1–6, <http://dx.doi.org/10.1007/s10482-017-0844-4>.